

Nonequilibrium conductance through a benzene molecule in the Kondo regime

L. Tosi,¹ P. Roura-Bas,² and A. A. Aligia¹

¹*Centro Atómico Bariloche and Instituto Balseiro,
Comisión Nacional de Energía Atómica, 8400 Bariloche, Argentina*

²*Dpto de Física, Centro Atómico Constituyentes,
Comisión Nacional de Energía Atómica, Buenos Aires, Argentina*

(Dated: March 8, 2013)

Starting from exact eigenstates for a symmetric ring, we derive a low-energy effective generalized Anderson Hamiltonian which contains two spin doublets with opposite momenta and a singlet for the neutral molecule. For benzene, the singlet (doublets) represent the ground state of the neutral (singly charged) molecule. We calculate the non-equilibrium conductance through a benzene molecule, doped with one electron or a hole (i.e. in the Kondo regime), and connected to two conducting leads at different positions. We solve the problem using the Keldysh formalism and the non-crossing approximation (NCA). When the leads are connected in the *para* position (at 180 degrees), the model is equivalent to the ordinary impurity Anderson model and its known properties are recovered. For other positions, there is a partial destructive interference in the cotunneling processes involving the two doublets and as a consequence, the Kondo temperature and the height and width of the central peak (for bias voltage V_b near zero) of the differential conductance $G = dI/dV_b$ (where I is the current) are reduced. In addition, two peaks at finite V_b appear. We study the position of these peaks, the temperature dependence of G and the spectral densities. Our formalism can also be applied to carbon nanotube quantum dots with intervalley mixing.

PACS numbers: 73.23.-b, 73.22.-f, 75.20.Hr

I. INTRODUCTION

Single molecule electronic devices are being extensively studied because they offer perspectives for further miniaturization of electronic circuits with important potential applications¹⁻⁵, and also due to its intrinsic interest in basic research, for example as realizations of the Kondo effect, which manifests experimentally by an increased conductance at low temperatures^{6,7}. In addition, quantum phase transitions involving partially Kondo screened spin-1 molecular states were induced changing externally controlled parameters⁸⁻¹⁰. These experiments could be explained semiquantitatively using extensions of the impurity Anderson model treated with either the numerical renormalization group (NRG)⁹⁻¹¹ or with the non-crossing approximation (NCA)¹⁰⁻¹³. This approximation allows calculations out of equilibrium and in particular at finite bias voltage.

The conductance through benzene-1,4-dithiol has been measured using a mechanically controllable break junction¹⁴. More recently, other molecules containing benzene rings or related phenyl groups were studied^{15,16}. The states nearer to the Fermi level in benzene are built from the 2p states of the C atoms which lie perpendicular to the plane of the molecule (the so called π states). Transport through aromatic molecules and in particular through benzene in different geometries has been calculated by several groups¹⁷⁻²¹. Hettler *et al.*¹⁷ started from an exact calculation of an extended Hubbard model for the π states, and included the coupling to the leads perturbatively (for couplings smaller than the temperature), to calculate the current through neutral benzene under an applied bias voltage V_b for zero gate voltage V_g , includ-

ing radiative relaxation. Cardamone *et al.*¹⁸, described the molecule by a Hubbard model supplemented by repulsions at larger distance (the so called Pariser-Parr-Pople (PPP) Hamiltonian²²) using the self-consistent Hartree-Fock approximation. The conductance of the effective one-body problem for small V_g has been calculated using the Landauer-Büttiker formalism²³. The authors propose to exploit the destructive interference and to control the conductance of the device introducing symmetry breaking perturbations. Similar results were obtained using *ab initio* calculations¹⁹, which have also the drawback of neglecting the effect of correlations.

This effect together with that interference were included in more recent works which started from the exact solution of the PPP model^{20,21}, as well as in previous works which discussed the conductance through molecules or rings of quantum dots (QDs) threaded by a magnetic flux²⁴⁻²⁹. The conductance G depends on which sites of the molecule are coupled to the leads and on interference phenomena related to the symmetry of the system^{20,21,29}. For certain conditions G can be totally suppressed and restored again by symmetry-breaking perturbations²¹. While the main conclusions are safe in general, the coupling to the leads was included in some perturbative approach, either by a Liouville equation method²⁰, or using the Jagla-Balseiro formula (JBF) for the conductance^{24,26}, which assumes a non-degenerate ground state. These approaches miss the Kondo effect which is non-perturbative in the coupling to the leads²⁸. This is particularly important when the gate voltage is such that the benzene molecule becomes charged, because the conductance is larger²¹.

The JBF has also been used to predict that the trans-

mittance integrated over a finite energy window^{24,25,27} and the equilibrium conductance^{25,29} through a ring of strongly correlated one-dimensional systems display dips as a function of the applied magnetic flux at *fractional* values of the flux quantum, due to destructive interference at crossings of levels with different charge and spin quantum numbers²⁹ (as a consequence of spin-charge separation). Recently, we have confirmed the presence of dips in the current under a finite applied bias voltage at low temperatures, using the low-energy effective Hamiltonian, consisting of two doublets and a singlet, for the case of perfect destructive interference³⁰. At equilibrium (zero bias voltage), the conductance as a function of temperature for this particular case has been calculated with NRG³¹. The interplay between interference and interactions were also studied for spinless electrons in multilevel systems³², and benzene attached to two leads³³. Although at first sight they might seem artificial, spinless models describe effective Hamiltonians for realistic systems under applied magnetic fields^{34,35}.

Interference phenomena were observed in systems with QDs. In a multilevel QD, the crossing of levels and the ensuing destructive interference has been induced applying magnetic field to a system with large, level-dependent g factors³⁴. Aharonov-Bohm oscillations were observed in systems involving two quantum dots (QDs)³⁶.

In this work, we construct the low-energy effective Hamiltonian for a ring of n sites with one orbital per site and symmetry that includes the point group C_{nv} (or C_n), weakly coupled to two conducting leads, retaining for n even, the lowest singlet with n particles and the two lowest doublets with $n + 1$ particles (electrons or holes depending on the sign of the applied gate voltage V_g). For n odd, the charge of doublets and singlets are interchanged. Using a gauge transformation, three of the four hopping matrix elements between the doublets and the leads can be made real. The phase of the fourth one ϕ is in general different from zero and depends on the position of the leads and the wave vectors of the states involved. While the coupling to the leads should be small in such a way that the neglected states do not affect the calculations³⁷, we treat this coupling in a self-consistent, non perturbative way, using the the Keldysh formalism of the NCA for problems out of equilibrium^{13,38,39}, appropriately extended for this problem. We calculate the non-equilibrium conductance for case of benzene ($n = 6$) in a regime of V_g for which the doublets are favored, leading to Kondo effect.

For the case of one doublet, comparison of NCA with NRG results⁴⁰, shows that the NCA describes accurately the Kondo physics. The leading behavior of the differential conductance for small voltage and temperature⁴¹ agrees with alternative Fermi-liquid approaches^{42,43}, and the temperature dependence of the conductance practically coincides with the NRG result over several decades of temperature⁴¹. A shortcoming of the NCA is that, at very low temperatures, it introduces an artificial spike at the Fermi energy in the spectral density when the ground

state of the system without coupling to the leads (zero hybridization) is non-degenerate, although the thermodynamic properties continue to be well described³⁸. This is not important in the present work, where the doublets are below the singlet and there is no applied magnetic field. Another limitation of the method is that it is restricted to temperatures above $\sim T_K/20$, where T_K is the Kondo temperature. In our case, this limitation is not important either because as we shall see, the conductance has already saturated at temperatures above this limit. A virtue of the NCA that becomes important in our case, is its ability to capture features at high energies, such as peaks in the spectral density out of the Fermi level, which might be lost in NRG calculations⁴⁴. An example is the plateau at intermediate temperatures observed in transport through C_{60} molecules for gate voltages for which triplet states are important^{8,10}, which was missed in early NRG studies, but captured by the NCA^{12,13}. More recent NRG calculations using tricks to improve the resolution⁴⁵, have confirmed this plateau¹⁰.

For the particular case of perfect destructive interference $\phi = \pi$ (which does not correspond to benzene), the spectral densities⁴⁶ and the non-equilibrium current³⁰, were calculated before for symmetric coupling to the leads. In this case, the model is equivalent to an SU(4) impurity Anderson model, used by several authors⁴⁷⁻⁵⁰ to interpret transport experiments in C nanotubes^{51,52}, and more recently in Si fin-type field effect transistors⁵³. If in addition a finite splitting δ between the doublets is allowed, the symmetry is reduced to SU(2). The transition between SU(4) and SU(2) has been studied by several authors using the NCA^{30,46,49,53}. In particular we found that the reduction of the Kondo temperature with δ agrees very well with a simple formula obtained from a variational calculation^{30,46}. An important difference between our model and those used for other systems is the connectivity with the leads, which imposes that in our case, the equilibrium conductance $G(V_b = 0)$ is zero in the SU(4) limit $\delta = 0$, and therefore, it is not proportional to the total spectral density of states³⁰. For $\phi = \pi$ and any δ , it is possible to calculate $G(0)$ in terms of static occupations, using a generalized Friedel-Langreth sum rule³⁰, due to the simplicity of the “orbital” field δ , which acts in the same way as a magnetic field, and allows to relate the phase shift for each channel and spin with the respective occupancies. In turn, these phase shifts enter a general Fermi-liquid expression for the conductance obtained by Pustilnik and Glazman⁵⁴. This is not possible for general ϕ . Therefore a novel treatment is needed to calculate the conductance even at equilibrium.

Recently, it has been shown that in carbon nanotube QDs with disorder induced valley mixing, the SU(4) symmetry is broken and the tunnel couplings to metallic leads become complex and depend on the applied magnetic field⁵⁵. The present formalism is also suitable to study this case.

The paper is organized as follows. In Section II we describe the construction of the effective Hamiltonian. The




FIG. 1. (Color online) Scheme of the relevant matrix elements at low energies.

approximations and the equation for the current are presented in Section III. Section IV contains the numerical results for transport through the benzene molecule and their interpretation. Section V contains a summary and a short discussion. In appendix A we show that the NCA conserves the current, extending the existing demonstration for the case of one doublet³⁸. Appendix B contains some numerical details.

II. MODEL

The effective model, which is represented at the right of Fig. 1, contains a singlet with total wave vector K_0 (usually 0 or π) and two doublets with wave vectors K_1 and K_2 , that represent the states of lowest energy of two neighboring configurations of a ring of n sites with symmetry C_{nv} . In the case of benzene, they correspond to the singlet ground state, invariant under rotations ($K_0 = 0$) and two degenerate doublets with total wave vectors $\pm K$, which are the ground state of the molecule for one added electron or hole, depending on the sign of the applied gate voltage^{20,21,29}. For one added hole $K = \pi/3$, while for one added electron $K = 2\pi/3$. The effective Hamiltonian is

$$H = E_s |s\rangle\langle s| + \sum_{i\sigma} E_i |i\sigma\rangle\langle i\sigma| + \sum_{\nu k\sigma} \epsilon_{\nu k} c_{\nu k\sigma}^\dagger c_{\nu k\sigma} + \sum_{i\nu k\sigma} (V_i^\nu |i\sigma\rangle\langle s| c_{\nu k\sigma} + \text{H.c.}), \quad (1)$$

where the singlet $|s\rangle$ and the two doublets $|i\sigma\rangle$ ($i = 1, 2$; $\sigma = \uparrow$ or $\downarrow$) denote the localized states, $c_{\nu k\sigma}^\dagger$ create conduction states in the left ($\nu = L$) or right ($\nu = R$) lead, and V_i^ν describe the hopping elements between the leads and both doublets, assumed independent of k . This hopping element is calculated as²⁹

$$V_i^\nu = t_\nu \langle i\sigma | c_{j_\nu\sigma}^\dagger | s \rangle, \quad (2)$$

where $c_{j\sigma}^\dagger$ creates an electron (or hole) with spin σ at the π orbital at site j (1 to n) of the molecule, j_ν denotes the site connected to the lead ν , and t_ν is the hopping between this site and the lead ν .

Choosing adequately the phases of the gauge transformation $|i\sigma\rangle \rightarrow e^{i\varphi_i} |i\sigma\rangle$, both V_i^L can be made real and

positive, and by reflection symmetry $V_1^L = V_2^L$. Using rotational symmetry, it is easy to see that²⁹

$$V_i^R = (t_R/t_L) V_i^L \exp[-i(j_R - j_L)(K_i - K_0)], \quad (3)$$

where K_i is the wave vector of $|i\sigma\rangle$. The phase of V_1^R can be absorbed by a gauge transformation in the $c_{Rk\sigma}$, rendering it real and positive. The remaining matrix element is $V_2^R = V_1^R e^{-i\phi}$, where $\phi = (j_R - j_L)(K_2 - K_1)$. This result is general for a ring geometry.

For benzene $K_2 - K_1 \equiv \pm 2\pi/3$. Then, if the leads are connected in the *para* position ($j_R - j_L = 3$), $\phi \equiv 0$, while in the *ortho* ($j_R - j_L = 1$) or *meta* ($j_R - j_L = 2$) positions, $\phi \equiv \pm 2\pi/3$. Note that the sign of ϕ does not affect our results. For simplicity we shall assume $|t_R/t_L| = 1$ in the calculations presented here. Then, the hopping of the leads to the relevant states of the benzene molecule are $V_1^L = V_2^L = V_1^R = V$, $V_2^R = V e^{-i\phi}$, with $\phi = 0$ for the *para* position, and $\phi = \pm 2\pi/3$ for the other two possibilities of connecting the leads.

For $\phi = 0$, the state $|B\sigma\rangle = (|1\sigma\rangle - |2\sigma\rangle)/\sqrt{2}$ decouples from the leads and the transport properties of the system are the same as those for a single level $[|A\sigma\rangle = (|1\sigma\rangle + |2\sigma\rangle)/\sqrt{2}]$ QD connected to the leads, with hopping $\sqrt{2}V$, which are well known^{38,56–59}. Another known limit of the model, which cannot be realized in benzene molecules, but in rings of a number of sites multiple of four²¹ is $\phi = \pi$. In this case, $|A\sigma\rangle$ ($|B\sigma\rangle$) is coupled only to the left (right) lead and the model is equivalent to an SU(4) Anderson model^{30,46}. However, in contrast to the case of C nanotubes^{47–50} and Si fin-type field effect transistors⁵³, the conductance vanishes identically as a consequence of perfect destructive interference^{30,31}. An important technical difference is that in our model, the hybridization matrices of the states $|i\sigma\rangle$ with the left and right leads are not proportional for $\phi \neq 0$. If the degeneracy between the levels can be lifted (for example applying an external flux²⁹), the symmetry is reduced to SU(2) and a finite conductance is restored^{30,31,46}. In C nanotubes with disorder induced valley mixing⁵⁵, the effective model becomes very similar to ours, including a finite level splitting and a phase ϕ , which depends on the applied magnetic field.

We shall use our previous results for the case $\phi = \pi$ and a finite level splitting $\delta = E_2 - E_1$ to help in the analysis of the results presented here for benzene.

III. THE FORMALISM

In this section we describe the extension of the non-crossing approximation (NCA) applied before for the Anderson model with infinite on-site repulsion out of equilibrium^{38,39}, to our effective Hamiltonian.

A. Representation of the Hamiltonian with slave particles

An auxiliary boson b , and four auxiliary fermions $f_{i\sigma}$ are introduced, so that the localized states are represented as

$$|s\rangle = b^\dagger|0\rangle, |i\sigma\rangle = f_{i\sigma}^\dagger|0\rangle, \quad (4)$$

where $|0\rangle$ is the vacuum. These pseudoparticles should satisfy the constraint

$$b^\dagger b + \sum_{i\sigma} f_{i\sigma}^\dagger f_{i\sigma} = 1. \quad (5)$$

Introducing it by a Lagrange multiplier, the effective Hamiltonian takes the form

$$H' = (E_s + \lambda)b^\dagger b + \sum_{i\sigma} (E_i + \lambda)f_{i\sigma}^\dagger f_{i\sigma} + \sum_{\nu k\sigma} \epsilon_{\nu k} c_{\nu k\sigma}^\dagger c_{\nu k\sigma} + \sum_{k\nu i\sigma} \left(V_i^\nu f_{i\sigma}^\dagger b c_{k\nu\sigma} + \text{H.c.} \right). \quad (6)$$

The NCA solves a system of self-consistent equations to obtain the Green functions of the pseudoparticles (described below), which is equivalent to sum an infinite series of diagrams (all those without crossings) in the corresponding perturbation series in the last term of H' , and afterwards project on the physical subspace of the constraint. The main new difficulties compared to the case of one doublet arise as a consequence of the matrix structure of the pseudofermion Green functions and self energies, which are absent in the SU(4) case or in this case with simple SU(2) symmetry breaking perturbations, as a magnetic⁵³ or "orbital"^{30,31,46} field.

B. Green functions

The lesser and greater Keldysh Green functions for the pseudoparticles for stationary non-equilibrium processes are defined as^{60,61}

$$\begin{aligned} G_{ij,\sigma}^<(t-t') &= +i\langle f_{j\sigma}^\dagger(t') f_{i\sigma}(t) \rangle, \\ D^<(t-t') &= -i\langle b^\dagger(t') b(t) \rangle, \\ G_{ij,\sigma}^>(t-t') &= -i\langle f_{i\sigma}(t) f_{j\sigma}^\dagger(t') \rangle, \\ D^>(t-t') &= -i\langle b(t) b^\dagger(t') \rangle, \end{aligned} \quad (7)$$

the retarded and advanced fermion Green functions are $G_{ij,\sigma}^r(t) = \theta(t)[G_{ij,\sigma}^>(t) + G_{ij,\sigma}^<(t)]$, $G_{ij,\sigma}^a = G_{ij,\sigma}^r + G_{ij,\sigma}^< - G_{ij,\sigma}^>$, and similarly for the boson Green functions replacing $G_{ij,\sigma}$ by D .

These Green functions correspond to the interacting (dressed) propagators and are determined selfconsistently within the NCA. Evaluating the corresponding diagrams in second order in the V_i^ν , and replacing the bare

propagators by the dressed ones, the expressions for the lesser self-energies take the form

$$\begin{aligned} \Pi^<(\omega) &= - \sum_{\nu lm\sigma} \Gamma_{lm}^\nu \int \frac{d\omega'}{2\pi} (1 - f_\nu(\omega' - \omega)) G_{ml,\sigma}^<(\omega'), \\ \Sigma_{lm,\sigma}^<(\omega) &= - \sum_{\nu} \Gamma_{lm}^\nu \int \frac{d\omega'}{2\pi} f_\nu(\omega - \omega') D^<(\omega'), \end{aligned} \quad (8)$$

where

$$\Gamma_{ij}^\nu(\omega) = 2\pi \sum_k V_i^\nu \bar{V}_j^\nu \delta(\omega - \epsilon_{\nu k}) \quad (9)$$

assumed independent of ω . Similarly, the greater self-energies become

$$\begin{aligned} \Sigma_{lm,\sigma}^>(\omega) &= \sum_{\nu} \Gamma_{lm}^\nu \int \frac{d\omega'}{2\pi} (1 - f_\nu(\omega - \omega')) D^>(\omega'), \\ \Pi^>(\omega) &= \sum_{\nu lm\sigma} \Gamma_{lm}^\nu \int \frac{d\omega'}{2\pi} f_\nu(\omega' - \omega) G_{ml,\sigma}^>(\omega'). \end{aligned} \quad (10)$$

As in the case of one doublet only³⁸, in the expressions for the retarded self energies, $\Sigma^r(t) = -\theta(t)(\Sigma^<(t) - \Sigma^>(t))$, $\Sigma^<$ can be neglected in comparison with $\Sigma^>$ due to the effect of the constraint. Then after Fourier transforming one obtains

$$\begin{aligned} \Sigma_{ij,\sigma}^r(\omega) &= i \int \frac{d\omega'}{2\pi} \frac{\Sigma_{ij,\sigma}^>(\omega')}{\omega - \omega' + i\eta}, \\ \Pi^r(\omega) &= i \int \frac{d\omega'}{2\pi} \frac{\Pi^>(\omega')}{\omega - \omega' + i\eta}, \end{aligned} \quad (11)$$

where η is a positive infinitesimal. The advanced self energies $\Sigma_{ij,\sigma}^a$ and Π^a are obtained changing the sign of η in the expressions above.

For the case of SU(4) symmetry or when this symmetry is broken by simple symmetry breaking fields^{30,31,46-50,53} ($\phi = \pi$ and symmetric coupling to the leads in our effective model), the fermion Green functions and self energies are diagonal in an appropriate basis and the self-consistency problem simplifies considerably. In the general case (including $\phi \equiv \pm 2\pi/3$ for benzene) one has to solve a matrix Dyson equation^{60,61} which includes not only the Keldysh (or +, - branch index in the Keldysh contour), but also the doublet index $i = 1, 2$ in the fermion case. For the retarded fermion Green functions and self energies, combining $G_{ij,\sigma}^r$ in a 2×2 matrix $\mathbf{G}^r$ and similarly for self energies and unperturbed Green functions $g_{ij,\sigma}^r = \delta_{ij}(\omega - \tilde{E}_i)^{-1}$, with $\tilde{E}_i = E - i + \lambda$ the Dyson equations take the simple form $\mathbf{G}^r = \mathbf{g}^r + \mathbf{g}^r \mathbf{\Sigma}^r \mathbf{G}^r$. Solving the system for the

$G_{ij,\sigma}^r$ we obtain

$$\begin{aligned} G_{11,\sigma}^r &= \frac{1}{\mathbb{D}}(\omega - \tilde{E}_2 - \Sigma_{22,\sigma}^r), \\ G_{12,\sigma}^r &= \frac{1}{\mathbb{D}}(\Sigma_{12,\sigma}^r), \\ G_{21,\sigma}^r &= \frac{1}{\mathbb{D}}(\Sigma_{21,\sigma}^r), \\ G_{22,\sigma}^r &= \frac{1}{\mathbb{D}}(\omega - \tilde{E}_1 - \Sigma_{11,\sigma}^r), \end{aligned} \quad (12)$$

where

$$\mathbb{D} = (\omega - \tilde{E}_1 - \Sigma_{11,\sigma}^r)(\omega - \tilde{E}_2 - \Sigma_{22,\sigma}^r) - \Sigma_{12,\sigma}^r \Sigma_{21,\sigma}^r.$$

For the boson, which has no subscripts, one has

$$D^r(\omega) = \frac{1}{\omega - \tilde{E}_s - \Pi^r}. \quad (13)$$

The advanced Green functions can be obtained from the retarded ones by the replacement $\eta \rightarrow -\eta$.

The remaining equations that relate the lesser and greater pseudoparticle Green functions with the corresponding self energies are (in compact matrix form for the fermions)⁶²

$$\begin{aligned} \mathbf{G}^{\lessgtr} &= \mathbf{G}^r \Sigma^{\lessgtr} \mathbf{G}^a, \\ D^{\lessgtr} &= D^r \Pi^{\lessgtr} D^a. \end{aligned} \quad (14)$$

We solve numerically the system of integral equations (8) to (14) for the pseudoparticle Green functions and self energies. The details are given in appendix B

For the calculation of the current, one needs the Green functions of the physical fermions $d_{i\sigma}^\dagger = |i\sigma\rangle\langle s| = f_{i\sigma}^\dagger b$. These functions, which we identify with the subscript $\mathbf{d}$, are defined in the same way as the pseudofermion ones [see Eqs. (7)], replacing $f_{i\sigma}$ by $d_{i\sigma}$. In terms of the auxiliary-particle Green functions, the lesser and greater physical Green functions take the form³⁸

$$\begin{aligned} G_{\mathbf{d}ij,\sigma}^<(\omega) &= i \int \frac{d\omega'}{2\pi Q} G_{ij,\sigma}^<(\omega' + \omega) D^>(\omega'), \\ G_{\mathbf{d}ij,\sigma}^>(\omega) &= i \int \frac{d\omega'}{2\pi Q} G_{ij,\sigma}^>(\omega' + \omega) D^<(\omega'), \end{aligned} \quad (15)$$

where $Q = \langle b^\dagger b + \sum_{i\sigma} f_{i\sigma}^\dagger f_{i\sigma} \rangle$ for a given Lagrange multiplier λ (see appendix B).

C. Equation for the current

Using general formulas for the current through a region with interacting electrons^{64,65} and the relation $G^r - G^a = G^> - G^<$ between Green functions, the current of our effective model for a spin degenerate system (without applied magnetic field) can be written as

$$\begin{aligned} I &= \pm \frac{ie}{h} \int d\omega \text{Tr}[(\mathbf{\Gamma}^L f_L(\omega) - \mathbf{\Gamma}^R f_R(\omega)) \mathbf{G}_{\mathbf{d}}^>(\omega) \\ &\quad + ((1 - f_L(\omega)) \mathbf{\Gamma}^L - (1 - f_R(\omega)) \mathbf{\Gamma}^R) \mathbf{G}_{\mathbf{d}}^<(\omega)], \end{aligned} \quad (16)$$

where the $+$ ($-$) sign correspond to the case in which the doublets have one more electron (hole) than the singlet, $f_\nu(\omega) = [\exp[(\omega - \mu_\nu)/kT] + 1]^{-1}$ where μ_ν is the chemical potential of the lead ν , and $\mathbf{\Gamma}^\nu$, $\mathbf{G}_{\mathbf{d}}^<$ and $\mathbf{G}_{\mathbf{d}}^>$ are 2×2 matrices with matrix elements given by Eqs (9) and (15). In particular, taking the unperturbed density of conduction states per spin $\rho = 1/(2D)$ where $2D$ is the band width, and symmetric coupling to the leads, we have

$$\begin{aligned} \mathbf{\Gamma}^L &= \frac{\pi V^2}{D} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \\ \mathbf{\Gamma}^R &= \frac{\pi V^2}{D} \begin{pmatrix} 1 & e^{i\phi} \\ e^{-i\phi} & 1 \end{pmatrix}. \end{aligned} \quad (17)$$

Note that unless $\phi = 0$, $\mathbf{\Gamma}^L$ is not proportional to $\mathbf{\Gamma}^R$. As a consequence, the trick used to relate the conductance at $V_b = 0$ with the spectral density of states^{64,65} cannot be used. We calculate the conductance $G = dI/dV_b$ by numerical differentiation of the current even for $V_b \rightarrow 0$. The traces appearing in Eq. (16) have the form

$$\begin{aligned} \text{Tr}(\mathbf{\Gamma}^R \mathbf{G}_{\mathbf{d}}^{\lessgtr}) &= \frac{\pi V^2}{D} [G_{11}^{\lessgtr} + G_{22}^{\lessgtr} + \cos(\phi)(G_{21}^{\lessgtr} + G_{12}^{\lessgtr}) \\ &\quad + i \sin(\phi)(G_{21}^{\lessgtr} - G_{12}^{\lessgtr})], \end{aligned} \quad (18)$$

and similarly for $\text{Tr}(\mathbf{\Gamma}^L \mathbf{G}_{\mathbf{d}}^{\lessgtr})$ replacing ϕ by 0.

Note that from the definition of the Green functions [see Eq. (7)], one realizes that the complex conjugates of the lesser and greater Green functions satisfy $\bar{G}_{ij}^{\lessgtr}(t) = -G_{ji}^{\lessgtr}(-t)$ and after Fourier transform $\bar{G}_{ij}^{\lessgtr}(\omega) = -G_{ji}^{\lessgtr}(\omega)$. Then, $G_{ii}^{\lessgtr}(\omega)$ and $G_{21}^{\lessgtr}(\omega) + G_{12}^{\lessgtr}(\omega)$ are pure imaginary, while $G_{21}^{\lessgtr}(\omega) - G_{12}^{\lessgtr}(\omega)$ is real.

In appendix A we show that the current from the left lead to the molecule equals that from the molecule to the right lead, so that the current is conserved within the NCA. Other approximations, like perturbation theory in the Coulomb repulsion out of the electron-hole symmetric point do not conserve the current⁴³, unless some tricks are used^{63,67,68}.

IV. NUMERICAL RESULTS

For the numerical calculations, we assume a constant density of states per spin of the leads ρ between $-D$ and D . We take the unit of energy as the total level width of both doublets: $\Gamma = \Gamma_{ii}^L + \Gamma_{ii}^R = 4\pi\rho V^2$, $i = 1, 2$. Also $\Gamma = 2\Delta$, where Δ , called the resonance level width, is half the width at half maximum of the spectral density of states of particles for each level in the non-interacting case. We restrict our present study to gate voltages such that $E_s - E_i \gg \Delta$, for which correlations play a more important role, the Kondo effect develops and the conductance is higher. Without loss of generality, we take $\epsilon_F = E_s = 0$, where ϵ_F is the Fermi level of

the leads without applied bias voltage. We define $E_d = E_1$, $\delta = E_2 - E_1$. For a bias voltage V_b , the chemical potentials are $\mu_L = eV_b/2$, $\mu_R = -eV_b/2$.

In the numerical results an important low-energy scale is the Kondo temperature T_K . It is known that for impurity Anderson models like ours for $\delta = 0$ in the Kondo regime, the spectral density shows one charge-transfer peak at energy E_d below the Fermi energy of total width of the order of NT for $SU(N)$ models^{46,47,66}. For finite on-site Coulomb repulsion U there is another charge-transfer peak at energy $E_d + U$, but this is shifted to infinite energies in our case. In addition, there is a narrow peak near the Fermi energy of width of the order of T_K . In this work we define T_K as the half width at half maximum of this peak. For finite δ additional peaks appear as shown in Section IV B.

A. Conductance out of equilibrium

In Fig. 2 we show the differential conductance $G = dI/dV_b$ as a function of bias voltage V_b , for the leads connected at 180 degrees (in the *para* position), for which $\phi = 0$, and in the other two positions ($\phi = \pm 2\pi/3$), for several values of the energy of the doublets $E_1 = E_2 = E_d$. The charge-transfer energy $\epsilon_F - E_d$ is the energy necessary to take a particle (electron or hole) from the Fermi energy, and add it to the localized singlet, forming a doublet, in absence of the hopping to the leads. It is tuned modifying the gate voltage V_g : E_d decreases with increasing V_g . The temperature in the curve was fixed at $T = 0.05T_K$. Since we have assumed symmetric coupling to both leads, $G(-V_b) = G(V_b)$ and only positive V_b are displayed in the figure.

For the *para* position, as explained in Section 2, the problem is equivalent to the transport through a single level quantum dot, and G shows a single peak centered near zero voltage and width of the order of $2kT_K/e$ ⁶⁹. We remind the reader that half the width of $G(V_b)$ for $T = 0$ times the electric charge e , half the width of the spectral density and the temperature (times the Boltzmann constant) T at which $G(T)$ for $V_b = 0$ falls to half of the maximum value, are all of the same order⁶⁹. This is natural from the expected universal behavior of the (single level) Anderson model in the Kondo regime, in which only one energy scale T_K survives. This Kondo temperature varies exponentially with the charge-transfer energy E_d , and as a consequence, the width of the central peak observed in the figure also decreases exponentially with increasing $|E_d|$, as observed in Fig. 2

When the molecule is attached to the leads at 60 or 120 degrees, three main changes occur in $G(V_b)$ in comparison to the previous case: i) The maximum of $G(0)$ is lower, ii) The width of the central peak is narrower and scales in a different way with E_d , and iii) two new peaks appear at finite V_b . i) is due to the effect of partial destructive interference between the current transmitted by the two doublets. In the *para* position, for which all




FIG. 2. (Color online) Differential conductance at low temperature through a benzene molecule as a function of bias voltage for several values of the energy of the doublets, with the leads connected in the *para* (red dashed line) and *ortho* or *meta* (full black line) positions.

hybridization to the leads have the same sign ($\phi = 0$), $G(0)$ equals the quantum of conductance $2e^2/h$ for symmetric coupling to the leads (as we have assumed in the present calculations). In the opposite case (hypothetical for benzene) of $\phi = \pi$, there is perfect destructive interference and G vanishes^{30,31}. Transport through the benzene molecule connected in the *ortho* or *meta* positions, correspond to an intermediate situation ($\phi = \pm 2\pi/3$), and therefore, an intermediate value of $G(0)$ is expected. For asymmetric coupling to the leads, as it is usually the case in transport through molecules in devices built by electromigration^{8,10}, $G(0)$ decreases strongly and it is not possible to distinguish between different positions of the leads connected to the benzene molecule from the maximum value of G . Note that this value decreases slightly




FIG. 3. (Color online) Differential conductance for $\phi = \pi$ and different values of the level splitting. Other parameters are $E_d = -4$, $T = 0.05T_K$.

with decreasing T_K . The feature, which is probably the most useful to distinguish experimentally among the different positions of the conducting leads is the presence of the side peaks in $G(V_b)$, which seem to depend weakly on the charge transfer energy $\epsilon_F - E_d$, in contrast to the width of the central peak.

In order to help in the interpretation of the results, we calculate $G(V_b)$ for $\phi = \pi$, but allowing for a finite splitting $\delta = E_2 - E_1$ between both doublets. δ acts as a symmetry breaking field on the SU(4) Kondo effect^{30,46}. The result is shown in Fig. 3. Clearly, $G(V_b)$ is qualitatively similar to the corresponding result for benzene with leads connected in the *ortho* or *meta* positions. The conductance, which vanishes in the SU(4) limit $\delta = 0$ is restored by a finite δ and two peaks at $eV_b = \pm\delta$ appear. This similarity suggest to interpret the results for benzene starting from those for $\phi = \pi$, $\delta = 0$ and thinking the difference between the coupling V_2^R for $\phi = \pm 2\pi/3$ and $\phi = \pi$ as a perturbation. This perturbation, in second order, introduces among other effects, an effective mixing between the levels, proportional to $V^2/|E_d|$, which leads to a splitting of the doublets. In fact, the position of the satellite peaks in Fig. 2 (ranging from 0.3 for $E_d = -3$ to 0.22 for $E_d = -5$) is roughly consistent with a $1/|E_d|$ dependence.

Note that in the case $\phi = 0$ in which this perturbation is expected to be the largest, only the symmetric combinations of left and right lead states hybridize with the impurity, while the antisymmetric ones remain decoupled. The spectral density of the latter is a delta function $\delta(\omega - E_d)$, without weight near the Fermi energy. This makes clear that actually the effective hopping of the two resulting split doublets is different and the proposed interpretation is rather qualitative.




FIG. 4. (Color online) Equilibrium spectral density for the benzene doublets as a function of frequency in the *ortho* and *meta* positions for different temperatures.




FIG. 5. (Color online) Total spectral density at equilibrium as a function of frequency for the effective model with $\phi = \pi$, $T = 0.05T_K$ and different values of the level splitting.

B. Spectral densities

As a further test of the interpretation outlined above, we compare the spectral density of states of both models. The spectral density

$$\rho_{i\sigma}(\omega) = (G_{\mathbf{d}i\sigma}(\omega + i\eta) - G_{\mathbf{d}i\sigma}(\omega - i\eta))/(2\pi i) \quad (19)$$

of both doublets for benzene connected to the leads in *ortho* or *meta* positions is represented in Fig. 4 for different temperatures and energies near the Fermi energy (the charge-transfer peak⁴⁶ is not shown). As T decreases below T_K , not only a peak develops near the Fermi energy, but also two side peaks (the most prominent for positive frequencies) are clearly present. In Fig. 5 we show the total spectral density for $\phi = \pi$, very low temperatures and different values of δ . In this case, the peak near to the Fermi energy corresponds to the lowest doublet and the peak at energy near δ corresponds to the highest doublet⁴⁶. In contrast, each peak in Fig. 4 is expected to come from some linear combination of both doublets which is not easy to identify. Another difference apparent in the figure is that the side peak at positive frequencies for benzene is sharper and more asymmetric




FIG. 6. (Color online) Conductance as a function of temperature for leads connected in the *ortho* or *meta* position and several values of the energy of the doublets.

than the corresponding one for $\phi = \pi$ and finite δ . This might be due in part to a smaller effective hybridization of the excited doublet in the case of benzene, leading to a narrower peak, as in the limit $\phi \rightarrow 0$ discussed above.

In spite of these differences, the spectral density for $\phi = \pi$ and finite δ shows the development of a satellite peak departing from the central one as δ increases. The spectral density for benzene shows the same qualitative features (a peak near the Fermi energy and another one at positive frequencies) which can be interpreted, in a qualitative first approximation, as coming from an effective δ . This peak at finite energy in turn gives rise to the side peaks in $G(V_b)$ (although the spectral densities are modified with the applied bias voltage and the peaks are blurred).

C. Temperature dependence of the conductance

In Fig. 6, we show the evolution with temperature of the conductance $G(T)$ for $V_b = 0$, several values of E_d , and conducting leads in the *ortho* or *meta* position. $G(T)$ was obtained differentiating the current, as explained in Section III C. In order to be able to display all curves in the same scale, a logarithmic scale of relative temperatures T/T_K is used, where the characteristic energy scale T_K is given by the half width at half maximum of the equilibrium spectral density $\rho_{i\sigma}(\omega)$. The resulting value of T_K is 3×10^{-2} , 3.6×10^{-3} and 4.8×10^{-4} for $E_d = -3$, -4 and -5 respectively. The peak at finite energy in the spectral density at low temperatures (which lies between 0.3 for $E_d = -3$ to 0.22 for $E_d = -5$) results in a structure in $G(T)$ at T/T_K near 10, 70 and 400 for $E_d = -3$, -4 and -5 respectively. While for $E_d = -3$ this structure is hidden in the main peak, a kind of plateau is evident for the other values of E_d at the corresponding positions, as shown in the inset of Fig. 6.

This structure is reminiscent of the plateau observed by transport experiments in the “triplet” side of the quantum phase transition in C_{60} ^{8,10}, and explained

by NCA^{10,12,13} and NRG calculations with improved resolution¹⁰. In our case, to represent accurately the high energy features, it was necessary to use a special mesh in the frequency axis with more points near the high energy peak in the pseudofermion spectral densities (see appendix B).

In previous work^{30,46}, we have found that for $\phi = \pi$ and any level splitting $\delta = E_2 - E_1$, T_K (except for a constant of order unity) is very accurately given by the expression

$$T_K = \{(D + \delta)D \exp[\pi E_1/(2\Delta)] + \delta^2/4\}^{1/2} - \delta/2, \quad (20)$$

obtained from a simple variational wave function that generalizes the proposal of Varma and Yafet⁷⁰ for the simplest impurity Anderson model. This equation interpolates between the SU(4) limit $\delta = 0$, for which T_K has the largest value $T_K^4 = D \exp[\pi E_1/(4\Delta)]$ and the one-level SU(2) limit $\delta \rightarrow +\infty$. The effect of δ on T_K is small when $\delta < T_K^4$, while for larger δ , T_K decreases strongly. Note that T_K^4 coincides with the Kondo temperature of the equivalent one-level effective Anderson for benzene connected in the *para* position (because the effective one-level resonant level width contains a factor 2, see Section II). Therefore, we can interpret the fact that the effective energy scale in the *ortho* and *meta* positions is smaller than that in the *para* positions (compare the widths of the central peaks in Fig. 2) as an effect of the effective level splitting caused by a phase ϕ different from π .

In other words, it seems that the position of the side peaks in $G(V_b)$ (or spectral densities) and the plateau in $G(T)$ indicate an effective δ , which decreases the characteristic energy scale T_K . While we do not expect Eq. (20) to describe accurately T_K using this effective δ , the fact that the difference between the energy scales for the *para* and the other positions is more noticeable for δ larger than T_K^4 (T_K for the *para* case), suggests that this equation might be useful for a qualitative understanding, and supports the interpretation of our results.

V. SUMMARY AND DISCUSSION

We have constructed the effective Hamiltonian for transport through a symmetric ring (point group including C_{nv} as a subgroup) with one orbital per site, including a singlet and two degenerate doublets of a neighboring configuration for the isolated ring. This includes for example the ground state of the isolated ring for one electron per site, and another with one added electron or hole. The resulting effective generalized Anderson Hamiltonian describes however more general situations. An extension to the case in which a magnetic flux is added (reducing the symmetry to C_n and breaking the doublet degeneracy) is trivial. Partial destructive interference occurs in general when two levels with N particles are near to another one with $N \pm 1$ particles²⁹. Therefore, the situation of one singlet and two doublets appears frequently.

From the wave vectors of these states, it is easy to calculate the phase ϕ of the effective model following the lines of Section II. Therefore, we expect that our formalism can be used in a variety of physically relevant systems.

We have used the resulting effective Hamiltonian to calculate the non-equilibrium transport through a benzene molecule with conducting leads connected in different positions, using an appropriate generalization of the NCA to include the Kondo effect as well as effects of partial destructive interference between the transport channels through both doublets. While for leads connected in the *para* position, the conductance G as a function of voltage and temperature is similar to that well known for the case of only one doublet, in the other positions the peak in $G(V_b)$ near zero bias voltage V_b is narrower and of smaller intensity as an indication of the relative position of the conducting leads. In addition, $G(V_b)$ displays two additional side peaks at finite $\pm V_b$, and a characteristic plateau is present in the conductance as a function of temperature $G(T)$. These finite energy features are probably easier to detect experimentally as an indication of the relative position of the conducting leads.

These results for the leads connected in the *ortho* and *meta* positions, which are due to partial destructive interference, can be interpreted starting from those with total destructive interference (corresponding to a phase $\phi = \pi$) for which the effective model has SU(4) symmetry, and adding a symmetry breaking splitting δ between effective doublets caused by the remaining term treated as a perturbation.

For our results to be valid, the coupling to the leads should be sufficiently small, so that excited states, not included in the effective Hamiltonian, do not play an important role. The effect of these states can be estimated for each particular case. See for example Ref.³⁷. The magnitude of the coupling can be controlled for example in break junctions^{6,9,14}.

Our effective model, including a finite level splitting δ from the beginning, and a general phase ϕ describes carbon nanotube QDs with disorder induced valley mixing⁵⁵. Therefore, our formalism supplemented by realistic calculations of δ and ϕ might be used to calculate the conductance of these systems.

In this work, we have neglected the effect of phonons, which can modify the effective parameters of the model^{72,73}, and also affect the interference phenomena⁷⁴. We have also neglected electrostatic interactions between the leads and the molecules. In the weak coupling regime, it has been shown that interaction with image charges for the doped molecules leads to a symmetry breaking and a splitting of the degenerate levels that also affects the interference phenomena⁷⁵. As we have shown, in the specific case of benzene, the coupling to the leads already originates a splitting of this kind. Therefore, we expect that these electrostatic effects are more relevant in the case of perfect destructive interference.

ACKNOWLEDGMENTS

We thank CONICET from Argentina for financial support. This work was partially supported by PIP 11220080101821 of CONICET and PICT R1776 of the ANPCyT, Argentina.

Appendix A: Current conservation

Here we present a brief demonstration that within NCA, the current established between the left metallic contact and the central region (ring) I_L , is the same as the current flowing from this region to the right metallic contact I_R . For simplicity we assume that there is no applied magnetic field. The demonstration can be easily extended to the case of inequivalent spins. Adding both spins the expressions of the currents are⁶⁴

$$I_L = \pm \frac{2ie}{h} \int d\omega \text{Tr}\{\Gamma^{\mathbf{L}}[f_L(\omega)\mathbf{G}_{\mathbf{d}}^>(\omega) + (1 - f_L(\omega))\mathbf{G}_{\mathbf{d}}^<(\omega)]\}, \quad (\text{A1})$$

$$I_R = \pm \frac{-2ie}{h} \int d\omega \text{Tr}\{\Gamma^{\mathbf{R}}[f_R(\omega)\mathbf{G}_{\mathbf{d}}^>(\omega) + (1 - f_R(\omega))\mathbf{G}_{\mathbf{d}}^<(\omega)]\}, \quad (\text{A2})$$

where the symbol $\pm$ and the meaning of the matrices is explained in Section III [actually Eq. (16) is the average of Eqs. (A1) and (A2)].

We use Eqs. (15) to replace the physical Green functions by their expressions in terms of the Green functions for the auxiliary particles. Denoting $\mathbf{G}$ the 2×2 matrix with the pseudo fermion Green functions for a given spin, one obtains

$$I_L = \mp \frac{2e}{h} \int d\omega \int \frac{d\omega'}{2\pi Q} f_L(\omega) D^<(\omega') \text{Tr}[\Gamma^{\mathbf{L}}\mathbf{G}^>(\omega' + \omega)] + (1 - f_L(\omega)) D^>(\omega') \text{Tr}[\Gamma^{\mathbf{L}}\mathbf{G}^<(\omega' + \omega)] \quad (\text{A3})$$

$$I_R = \pm \frac{2e}{h} \int d\omega \int \frac{d\omega'}{2\pi Q} f_R(\omega) D^<(\omega') \text{Tr}[\Gamma^{\mathbf{R}}\mathbf{G}^>(\omega' + \omega)] + (1 - f_R(\omega)) D^>(\omega') \text{Tr}[\Gamma^{\mathbf{R}}\mathbf{G}^<(\omega' + \omega)]. \quad (\text{A4})$$

Using Eqs. (10) for the boson self-energies, and some algebra, the difference between both expressions becomes

$$I_L - I_R = \frac{-e}{h} \int \frac{d\omega'}{Q} [D^<(\omega')\Pi^>(\omega') - D^>(\omega')\Pi^<(\omega')], \quad (\text{A5})$$

which is easily seen to vanish using $D^{\lessgtr} = D^r \Pi^{\lessgtr} D^a$ [see Eqs. (14)].

Appendix B: Numerical procedure

In this appendix we give some details about the numerical solution of the system of integral equations.

We first solve the system that determines the retarded and greater Green functions and self energies for the auxiliary particles, Eqs. (10) to (14). After the system converges, we solve self-consistently Eqs. (8) and (14) for the lesser quantities.

Due to the gauge symmetry of the Hamiltonian in the representation of the auxiliary particles, $f_{i\sigma} \rightarrow \exp(i\lambda_0 t)f_{i\sigma}$, $b \rightarrow f_{i\sigma} \exp(i\lambda_0 t)b$, the Lagrange multiplier λ in Eq. (15) can be shifted by an arbitrary number λ_0 ³⁹. This is equivalent to a shift in the energies of the slave particles or of the frequency as $\omega \rightarrow \omega + \lambda_0$. Therefore, we can either set $Q = \langle b^\dagger b + \sum_{i\sigma} f_{i\sigma}^\dagger f_{i\sigma} \rangle = 1$ and calculate λ at each iteration or fix λ , calculate Q and divide

by Q in the evaluation of physical properties^{38,39}.

In the present problem, the spectral functions of the pseudo-fermions have two different peaks in contrast to the single one present in the ordinary Anderson model. We found that fixing λ as the solution of the equation $\text{Re}D(\omega, \lambda) = 0$ at each iteration, the position of the lowest peak of the auxiliary fermion spectral densities coincide with the zero of energy (the Fermi level) With this choice of λ , in order to make the numerical evaluations efficiently, we use a non equidistant frequency mesh ω_i with two different regions logarithmically spaced. One of them is aimed at the zero of energy $\omega = 0$ and the other one is centered near to the position of the second peak of the fermion spectral functions, $\omega = \delta$. Finally, to generate the non equidistant mesh we follow reference³⁹.

-
- ¹ Nitzan A and Ratner M A, 2003 *Science* **300** 1384
 - ² Venkataraman L, Klare J E, Nuckolls C, Hybertsen M S and Steigerwald M L, 2006 *Nature (London)* **442** 904
 - ³ Galperin M, Ratner M A, Nitzan A and Troisi A, 2008 *Science* **319** 1056
 - ⁴ van der Molen S J and Liljeroth P, 2010 *J. Phys. Condens. Matter* **22** 133001
 - ⁵ Cuevas J C and Scheer E, 2010 *Molecular Electronics: An Introduction to Theory and Experiment* (World Scientific, Singapore)
 - ⁶ Park J, Pasupathy A N, Goldsmith J I, Chang C, Yaish Y, Petta J R, Rinkoski M, Sethna J P, Abruña H D, P L McEuen and Ralph D C, 2002 *Nature (London)* **417** 722
 - ⁷ Lian W, Shores M P, Bockrath M, Long J R and Park H, *Nature (London)* 2002 **417** 725
 - ⁸ Roch N, Florens S, Bouchiat V, Wernsdorfer W and Balestro F, 2008 *Nature (London)* **453** 633
 - ⁹ Parks J J, Champagne A R, Costi T A, Shum W W, Pasupathy A N, Neuscamman E, Flores-Torres S, Cornaglia P S, Aligia A A, Balseiro C A, Chan G K-L, Abruña H D, and Ralph D C, 2010 *Science* **328** 1370
 - ¹⁰ Florens S, Freyn A, Roch N, Wernsdorfer W, Balestro F, Roura-Bas P and Aligia A A 2011 *J. Phys.: Condens. Matter* **23** 243202
 - ¹¹ Cornaglia P S, Roura-Bas P, Aligia A A and Balseiro C A 2011 *Europhys. Lett.* **93** 47005
 - ¹² Roura-Bas P and Aligia A A 2009 *Phys. Rev. B* **80** 035308
 - ¹³ Roura-Bas P and Aligia A A 2010 *J. Phys.: Condens. Matter* **22** 025602
 - ¹⁴ Reed M A, Zhou C, Muller C J, Burgin T P and Tour J M, *Science* **278** 252 (1997).
 - ¹⁵ Danilov A V, Kubatkin S, Kafanov S, Hedegård P, Stühr-Hansen N, Moth-Poulsen K, and Bjørnholm T, 2008 *Nano Lett.* **8** 1
 - ¹⁶ Dadosh T, Gordin Y, Krahne R, Khivrich I, Mahalu D, Frydman V, Sperling J, Yacoby A, and Bar-Joseph I, 2005 *Nature (London)* **436** 677
 - ¹⁷ Hettler M H, Wenzel W, Wegewijs M R, and Schoeller H, 2003 *Phys. Rev. Lett.* **90** 076805
 - ¹⁸ Cardamone D, Stafford C A, and Mazumdar S, 2006 *Nano Lett.* **6** 2422
 - ¹⁹ Ke S-H, Yang W, and Baranger H U, 2008 *Nano Lett.* **8** 3257
 - ²⁰ Begemann G, Darau D, Donarini A, and Grifoni M, 2008 *Phys. Rev. B* **77** 201406(R)
 - ²¹ Rincón J, Hallberg K, Aligia A A and Ramasesha S, 2009 *Phys. Rev. Lett.* **103**, 266807
 - ²² Pariser R and Parr R, 1953 *J. Chem. Phys.* **21** 466; Pople J A, 1953 *Trans. Faraday Soc.* **49** 1375 .
 - ²³ Büttiker M, 1986 *Phys. Rev. Lett.* **57** 1761.
 - ²⁴ Jagla E A and Balseiro C A, 1993 *Phys. Rev. Lett.* **70** 639
 - ²⁵ Hallberg K, Aligia A A, Kampf A P and Normand B, 2004 *Phys. Rev. Lett.* **93** 067203
 - ²⁶ Aligia A A, Hallberg K, Normand B and Kampf A P, 2004 *Phys. Rev. Lett.* **93** 076801
 - ²⁷ Friederich S and Meden V, 2008 *Phys. Rev. B* **77** 195122
 - ²⁸ Lobos A M and Aligia A A, 2008 *Phys. Rev. Lett.* **100** 016803; 2009 *Physica B* **404**, 3306
 - ²⁹ Rincón J, Aligia A A, and Hallberg K 2009 *Phys. Rev. B* **79** 035112
 - ³⁰ Roura-Bas P, Tosi L, Aligia A A, and Hallberg K, 2011 *Phys. Rev. B* **84**, 073406
 - ³¹ Izumida W, Sakai O, and Shimizu Y, 1998 *J. Phys. Soc. Jpn.* **67**, 2444
 - ³² Meden V and Marquardt F, 2006 *Phys. Rev. Lett.* **96** 146801
 - ³³ Bohr D and Schmitteckert P, 2012 *Ann. Phys.* **524**, 199
 - ³⁴ Nilsson H A, Karlström O, Larsson M, Caroff P, Pedersen J N, Samuelson L, Wacker A, Wernersson L-E and Xu H Q, 2010 *Phys. Rev. Lett.* **104** 186804
 - ³⁵ Tosi L and Aligia A A, 2011 *Phys. Status Solidi B* **248** 732
 - ³⁶ Hatano T, Kubo T, Tokura Y, Amaha S, Teraoka S, and Tarucha S, 2011 *Phys. Rev. Lett.* **106**, 076801
 - ³⁷ Lobos A M and Aligia A A, 2006 *Phys. Rev. B* **74** 165417
 - ³⁸ Wingreen N S and Meir Y, 1994 *Phys. Rev. B* **49** 11040
 - ³⁹ Hettler M H, Kroha J and Hershfield S, 1998 *Phys. Rev. B* **58** 5649
 - ⁴⁰ Costi T A, Kroha J and Wölffe P, 1996 *Phys. Rev. B* **53** 1850
 - ⁴¹ Roura-Bas P, 2010 *Phys. Rev. B* **81**, 155327
 - ⁴² Oguri A, 2005 *J. Phys. Soc. Jpn.* **74** 110
 - ⁴³ Aligia A A, 2012 *J. Phys. Condens. Matter* **24** 015306 ; references therein.
 - ⁴⁴ Vaugier L, Aligia A A and Lobos A M, 2007 *Phys. Rev. B* **76** 165112
 - ⁴⁵ Freyn A and Florens S, 2011 *Phys. Rev. Lett.* **107** 017201

- ⁴⁶ Tosi L, Roura-Bas P, and Aligia A A, 2012 *Physica B* **407** 3259
- ⁴⁷ Lim J S, Choi M-S, Choi M Y, López R and Aguado R, 2006 *Phys. Rev. B* **74** 205119
- ⁴⁸ Anders F B, Logan D E, Galpin M R, and Finkelstein G, 2008 *Phys. Rev. Lett.* **100** 086809
- ⁴⁹ Lipinski S and Krychowski D, 2010 *Phys. Rev. B* **81** 115327
- ⁵⁰ Büsser C A, Vernek E, Orellana P, Lara G A, Kim E H, Feiguin A E, Anda E V and Martins G B, 2011 *Phys. Rev. B* **83** 125404
- ⁵¹ Jarillo-Herrero P, Kong J, van der Zant H S J, Dekker C, Kouwenhoven L P and De Franceschi S, 2005 *Nature (London)* **434** 484
- ⁵² Makarovski A, Liu J, and Finkelstein G, 2007 *Phys. Rev. Lett.* **99** 066801
- ⁵³ Tettamanzi G C, Verduijn J, Lansbergen G P, Blaauboer M, Calderón M J, Aguado R, and Rogge S, 2012 *Phys. Rev. Lett.* **108** 046803
- ⁵⁴ Pustilnik M and Glazman L I, 2001 *Phys. Rev. Lett.* **87**, 216601
- ⁵⁵ Grove-Rasmussen K, Grap S, Paaske J, Flensberg K, Andergassen S, Meden V, Jorgensen H I, Muraki K, and Fujisawa T, 2012 *Phys. Rev. Lett.* **108**, 176802
- ⁵⁶ Izumida W, Sakai O, and Suzuki S, 2001 *J. Phys. Soc. Jpn.* **70** 1045
- ⁵⁷ Costi T A, 2001 *Phys. Rev. B* **64** 241310(R)
- ⁵⁸ Aligia A A and Proetto C R 2002 *Phys. Rev. B* **65** 165305
- ⁵⁹ Costi T A and Zlatić V, 2010 *Phys. Rev. B* **81**, 235127
- ⁶⁰ Lifshitz E M and Pitaevskii A L, 1981 *Physical Kinetics* (Pergamon, Oxford)
- ⁶¹ Mahan G D, 2000 *Many Particle Physics* (Kluwer/Plenum, New York)
- ⁶² In the general case, Eqs. (14) contain an additional term of the form $|G^r/g^r|^2 g^{\lessgtr}$ (see for example Eq. (7) of Ref.⁶³), but they vanish in this case because $(g_{ij,\sigma}^r)^{-1} = \delta_{ij}(\omega - \tilde{E}_i)$ and $g^{\lessgtr} \sim \delta_{ij}\delta(\omega - \tilde{E}_i)$.
- ⁶³ Aligia A A, 2006 *Phys. Rev. B* **74** 155125
- ⁶⁴ Meir Y and Wingreen N S 1992 *Phys. Rev. Lett.* **68** 2512
- ⁶⁵ Jauho A P, Wingreen N S, and Meir Y, 1994 *Phys. Rev. B* **50**, 5528
- ⁶⁶ Logan D E, Eastwood M P and Tusch M A, 1998 *J. Phys.: Condens. Matter* **10** 2673
- ⁶⁷ Levy Yeyati A, Martin-Rodero A, and Flores F, 1993 *Phys. Rev. Lett.* **71** 2991
- ⁶⁸ Monreal R C, Flores F, and Martin-Rodero A, 2010 *Phys. Rev. B* **82** 235412
- ⁶⁹ Tosi L, Roura-Bas P, Llois A M, and Aligia A A, 2012 *Physica B* **407** 3263
- ⁷⁰ Varma C M and Yafet Y, 1976 *Phys. Rev. B* **13** 2950
- ⁷¹ Langreth D C 1966 *Phys. Rev.* **150** 516
- ⁷² Härtle R and Thoss M, 2011 *Phys. Rev. B* **83** 115414
- ⁷³ Cornaglia P S, Usaj G, and Balseiro C A, 2007 *Phys. Rev. B* **76** 241403(R)
- ⁷⁴ Härtle R, Butzin M, Rubio-Pons O, and Thoss M, 2011 *Phys. Rev. Lett.* **107** 046802
- ⁷⁵ Kaasbjerg K and Flensberg K, 2011 *Phys. Rev. B* **84** 115457